arXiv:1511.06465v2 [math.FA] 23 Nov 2015

SOBOLEV FUNCTIONS IN THE CRITICAL CASE ARE UNIFORMLY CONTINUOUS IN s-AHLFORS REGULAR METRIC SPACES WHEN $s \leq 1$

XIAODAN ZHOU

Abstract. We prove that functions in the Hajłasz-Sobolev space $M^{1,s}$ on an s-Ahlfors regular metric space are uniformly continuous when $s \leq 1$.

1. Introduction

Sobolev functions $u \in W^{1,1}([a,b])$ are absolutely continuous. When $n \geq 2$, $u(x) = \log|\log|x||$ provides an example of a discontinuous Sobolev function in $W^{1,n}(B^n(0,e^{-1}))$. The aim of this paper is to investigate continuity of the Sobolev functions in the critical case, i.e. $p = n$ on more general metric spaces including fractals.

Definition 1. *Let (X,d,μ) be a metric space equipped with a Borel measure μ and let $s > 0$. We say that a metric measure space (X,d,μ) is s-Ahlfors regular if there is a constant $C_A \geq 1$, such that*

$$C_A^{-1} r^s \leq \mu(B(x,r)) \leq C_A r^s, \quad \text{for} \quad x \in X \quad \text{and} \quad 0 < r < \operatorname{diam}(X).$$

An important class of Ahlfors regular spaces is provided by fractals. Hutchinson [15], proved that fractals generated by iterated function systems satisfying the open set condition are Ahlfors regular.

If X is s-Ahlfors regular with respect to μ, then μ is comparable to the Hausdorff measure $\mathcal{H}^s$. So X is s-Ahlfors regular with respect to $\mathcal{H}^s$ [11, Exercise 8.11.].

Analysis on fractals has developed a great deal since eighties [3, 5, 17, 18, 25] and the theory of Sobolev spaces on metric spaces play an important role in this development. There are various extensions of the theory of Sobolev spaces to metric-measure spaces [1, 2, 4, 8, 9, 10, 11, 12, 24]. While the approach based on upper gradients (Cheeger and Newtonian Sobolev spaces) requires the space to be highly connected, it does not apply to fractals with limited connectivity properties. On the other hand, the theory of Hajłasz-Sobolev spaces is rich without any assumption on connectivity of the space, and Hajłasz-Sobolev spaces on fractals have been investigated in [13, 14, 22, 23, 26].

2010 *Mathematics Subject Classification.* 46E35, 28A80.
Key words and phrases. Sobolev functions, uniform continuity, Fractals, Ahlfors regular.
The author was partially supported by NSF grant DMS-1500647 of Piotr Hajłasz.

Definition 2. *Let (X, d, μ) be a metric space equipped with a Borel measure μ. For $0 < p < \infty$ we define the Hajłasz-Sobolev space $M^{1,p}(X, d, \mu)$ to be the set of all functions $u \in L^p(X)$ for which there exists nonnegative Borel functions $g \in L^p(X)$ such that*

$$|u(x) - u(y)| \leq d(x, y)\big(g(x) + g(y)\big) \quad \mu - a.e.$$

Denote by $D(u)$ the class of all nonnegative Borel functions g that satisfy the above inequality. Thus $u \in M^{1,p}(X, d, \mu)$ if and only if $u \in L^p(X)$ and $D(u) \cap L^p \neq \emptyset$. The space $M^{1,p}(X, d, \mu)$ is linear and we equip it with

$$||u||_{M^{1,p}} = ||u||_{L^p} + \inf_{g \in D(u)} ||g||_{L^p}.$$

Remark 1. If $p \geq 1$, $|| \cdot ||_{M^{1,p}}$ is a norm and $M^{1,p}$ is a Banach space [8]. For applications of Hajłasz-Sobolev spaces with $0 < p < 1$, see for example [9, 16, 19, 20, 21, 28].

The following result is the main theorem of this paper.

Theorem 1. *Let $(X, d, \mathcal{H}^s)$ be an s-Ahlfors regular metric space and $0 < s \leq 1$. If $u \in M^{1,s}(X, d, \mathcal{H}^s)$, then u is uniformly continuous. Moreover, there exists a constant $C > 0$, such that for any ball $B \subset X$,*

$$\operatorname*{osc}_B |u| = \sup_{x,y \in B} |u(x) - u(y)| \leq C\Big(\int_{2B} g^s d\mathcal{H}^s \Big)^{\frac{1}{s}}, \qquad (*)$$

where $g \in D(u) \cap L^s(X)$.

Here and in what follows by $2B$ we denote a ball concentric with B and with twice the radius.

Remark 2. For example, the ternary Cantor set C equipped with the Euclidean distance is s-Ahlfors regular with $s = \frac{\log 2}{\log 3}$, [6, Example 2.7].

Remark 3. The Sobolev functions in $W^{1,1}([a, b])$ are absolutely continuous. More generally, if the space X is compact, connected and $\mathcal{H}^1(X) < \infty$, the functions in the Newtonian Sobolev spaces $N^{1,1}(X, d, \mu)$ are also absolutely continuous [27].

Remark 4. The restriction that $s \leq 1$ is necessary in our statement, since we apply the reverse Minkowski inequality in the proof. We conjecture that there alway exists discontinuous Sobolev functions in $M^{1,s}(X)$ when X is an s-Ahlfors regular space with $s > 1$.

2. Proof of the main theorem

Let C denote a general constant whose value can change even in the same string of estimates.

Lemma 2. *Let X be an s-Ahlfors regular space. Then there is a constant $0 < C_0 < 1$ such that for any ball $B(x,r)$ with $0 < r < \operatorname{diam}(X)$, there is a sequence of balls $\{B_i\}_{i=1}^{\infty} = \{B(x_i, C_0^i r)\}_{i=1}^{\infty} \subset B(x,r)$ satisfying*

(1) *$B_i \cap B_j = \emptyset$, if $i \neq j$,*

(2) *$B(x_i, C_0^i r) \subset B(x, C_0^{i-1} r) \setminus B(x, C_0^i r)$.*

Proof. Since X is s-Ahlfors regular, there is a constant $C_A \geq 1$, such that for any ball $B(x,r) \subset X$ with $0 < r < \operatorname{diam}(X)$, we have

$$C_A^{-1} r^s \leq \mathcal{H}^s(B(x,r)) \leq C_A r^s.$$

We claim that for any $0 < c < C_A^{-2/s}$,

$$B(x,r) \setminus B(x,cr) \neq \emptyset.$$

For if not, $B(x,r) \setminus B(x,cr) = \emptyset$ would imply

$$C_A^{-1} r^s \leq \mathcal{H}^s(B(x,r)) = \mathcal{H}^s(B(x,cr)) \leq C_A (cr)^s$$

so $C_A^{-2/s} \leq c$, which is a contradiction. Fix $0 < c < C_A^{-2/s}$ and let

$$C_1 = \min\{\frac{c}{2}, \frac{\sqrt{5}-1}{2}\}.$$

Clearly, $0 < C_1 < 1$.

Fix a ball $B(x,r) \subset X$ with $0 < r < \operatorname{diam}(X)$ and let $y \in B(x,r) \setminus B(x,cr)$. Then

$$B(y, C_1 r) \subset B(x, C_1^{-1} r) \setminus B(x, C_1 r).$$

We can construct a sequence of balls $B_1, B_2 \ldots \subset B(x,r)$ in the following way. Let $\hat{r}_i = C_1^{2i-1} r$ and $\hat{B}_i = B(x, \hat{r}_i)$. By the previous argument, for each $\hat{B}_i$, there exists x_i such that

$$B(x_i, C_1 \hat{r}_i) \subset B(x, C_1^{-1} \hat{r}_i) \setminus B(x, C_1 \hat{r}_i).$$

Let $r_i = C_1 \hat{r}_i = C_1^{2i} r$, so

$$B(x_i, r_i) \subset B(x, r_{i-1}) \setminus B(x, r_i).$$

Let $C_0 = C_1^2$. Clearly, $0 < C_0 < 1$. Let $B_i = B(x_i, r_i) = B(x_i, C_0^i r)$. Since these balls are contained in disjoint annuli, $B_i \cap B_j = \emptyset$ for $i \neq j$.

From the construction, the balls in this sequence satisfy

$$B_i \subset B(x, C_0^{i-1} r) \setminus B(x, C_0^i r).$$

This completes the proof. □

Now, we can complete the proof of the main result follwoing some ideas from [7].

Proof. By definition, for $u \in M^{1,s}(X, d, \mathcal{H}^s)$, there exists $g \in L^s(X)$ such that

$$|u(x) - u(y)| \le d(x, y)\big(g(x) + g(y)\big),$$

when $x, y \in X \setminus E$ and $\mathcal{H}^s(E) = 0$.

Let $E_0 = \{x \in X \setminus E : g(x) < \infty\}$. Clearly, $\mathcal{H}^s(X \setminus E_0) = 0$. Fix a ball $B(z, r) \subset X$. We will prove that

$$\sup_{x,y \in B(z,r) \cap E_0} |u(x) - u(y)| \le C\Big(\int_{B(z,2r)} g^s d\mathcal{H}^s \Big)^{\frac{1}{s}}.$$

Let $x, y \in B(z, r) \cap E_0$. According to Lemma 2, there exists a sequence of disjoint balls $\{B_i\}_{i=1}^{\infty} = \{B(x_i, C_0^i r)\}_{i=1}^{\infty} \subset B(x, r)$ such that

$$B_i = B(x_i, C_0^i r) \subset B(x, C_0^{i-1} r) \setminus B(x, C_0^i r).$$

Let $r_i = C_0^i r$, then

$$B_i = B(x_i, r_i) \subset B(x, C_0^{-1} r_i) \setminus B(x, r_i).$$

Clearly, we can find $z_i \in B_i \cap E_0$ such that

$$\begin{aligned} g(z_i) \le \Big(\fint_{B_i} g^s d\mathcal{H}^s \Big)^{1/s} &= \Big(\frac{1}{\mathcal{H}^s(B_i)} \Big)^{1/s} \Big(\int_{B_i} g^s d\mathcal{H}^s \Big)^{1/s} \\ &\le \frac{C}{r_i} \Big(\int_{B_i} g^s d\mathcal{H}^s \Big)^{1/s}. \end{aligned}$$

Here the barred integral denotes the integral average.

Notice that $z_i \in B_i \subset B(x, C_0^{-1} r_i)$ implies that $d(x, z_i) \le C_0^{-1} r_i$. It follows that

$$\begin{aligned} |u(x) - u(z_i)| &\le d(x, z_i)\Big(g(x) + g(z_i)) \Big) \\ &\le C r_i g(x) + C \Big(\int_{B_i} g^s d\mathcal{H}^s \Big)^{1/s}. \end{aligned}$$

Hence, $\lim_{i \to \infty} u(z_i) = u(x)$. Thus,

$$\begin{aligned} |u(z_1) - u(x)| &\le \sum_{i=1}^{\infty} |u(z_i) - u(z_{i+1})| \\ &= \sum_{i=0}^{\infty} d(z_i, z_{i+1})\Big(g(z_i) + g(z_{i+1}) \Big). \end{aligned}$$

Since $z_i \in B_i$, $z_{i+1} \in B_{i+1}$ and $B_i, B_{i+1} \subset B(x, C_0^{-1} r_i)$, we have that $d(z_i, z_{i+1}) \leq Cr_i$. Since also $r_{i+1} = C_0 r_i$, the above inequality implies that

$$\begin{aligned} |u(z_1) - u(x)| &\leq \sum_{i=0}^{\infty} Cr_i \Big(\frac{1}{r_i} \big(\int_{B_i} g^s d\mathcal{H}^s \big)^{1/s} + \frac{1}{r_{i+1}} \big(\int_{B_{i+1}} g^s d\mathcal{H}^s \big)^{1/s} \Big) \\ &\leq C \sum_{i=0}^{\infty} \Big(\int_{B_i} g^s d\mathcal{H}^s \Big)^{1/s} \\ &\leq C \Big(\int_{B(x,r)} g^s d\mathcal{H}^s \Big)^{1/s}. \end{aligned} \tag{1}$$

The last inequality follows from the reverse Minkowski inequality and the fact that the balls in the sequence $\{B_i\}_{i=1}^{\infty} \subset B(x, r)$ are pairwise disjoint.

Notice that $z_1 \in B_1 \subset B(x, r) \subset B(z, 2r)$ and

$$g(z_1) \leq \frac{C}{r_1} \Big(\int_{B_1} g^s d\mathcal{H}^s \Big)^{1/s} \leq \frac{C}{r} \Big(\int_{B(z,2r)} g^s d\mathcal{H}^s \Big)^{1/s}.$$

Similarly, we can find $w_1 \in B(y, r) \subset B(z, 2r)$ such that

$$|u(w_1) - u(y)| \leq C \Big(\int_{B(y,r)} g^s d\mathcal{H}^s \Big)^{1/s}, \tag{2}$$

and

$$g(w_1) \leq \frac{C}{r} \Big(\int_{B(z,2r)} g^s d\mathcal{H}^s \Big)^{1/s}.$$

Since $z_1, w_1 \in B(z, 2r)$, we also have

$$\begin{aligned} |u(z_1) - u(w_1))| &\leq d(z_1, w_1)(g(z_1) + g(w_1)) \\ &\leq C \big(\int_{B(z,2r)} g^s d\mathcal{H}^s \big)^{1/s}. \end{aligned} \tag{3}$$

Now the inequalities (1), (2), (3) readily give

$$\sup_{x,y \in B(z,r) \cap E_0} |u(x) - u(y)| \leq C \Big(\int_{B(z,2r)} g^s d\mathcal{H}^s \Big)^{\frac{1}{s}}.$$

This and the absolute continuity of the integral imply uniform continuity of $u|_{X \setminus E_0}$. Hence u uniquely extends to a uniformly continuous function on X and the inequality $(*)$ follows. The proof is complete.

□

Acknowledgments

The author thanks Professor P. Hajłasz for bringing this topic to her attention and valuable suggestions to improve the paper. She also thanks Professor N. Shanmugalingam for her interests and helpful comments.

References

[1] L. Ambrosio, N. Gigli, G. Savar, *Gradient flows in metric spaces and in the space of probability measures*. Second edition. Lectures in Mathematics ETH Zrich. Birkhuser Verlag, Basel, 2008.

[2] L. Ambrosio and P. Tilli: *Topics on analysis in metric spaces*. Oxford Lecture Series in Mathematics and its Applications, 25. Oxford University Press, Oxford, 2004.

[3] M. Barlow: *Diffusions on fractals*. Lectures on probability theory and statistics (Saint-Flour, 1995), 1-121, Lecture Notes in Math., 1690, Springer, Berlin, 1998.

[4] J. Cheeger: Differentiability of Lipschitz functions on metric measure spaces. *Geom. Funct. Anal.*, 9(3):428–517, 1999.

[5] G. David and S. Semmes: *Fractured fractals and broken dreams: Self-similar geometry through metric and measure*. Oxford University Press, New York, 1997.

[6] K. Falconer: *Fractal geometry. Mathematical foundations and applications.* John Wiley & Sons, Inc., Hoboken, NJ, 2nd ed, 2003.

[7] P. Hajłasz: Geometric approach to Sobolev spaces and badly degenerated elliptic equations. *Nonlinear analysis and applications (Warsaw, 1994)*, 141–168, 1995.

[8] P. Hajłasz: Sobolev spaces on an arbitrary metric space. *Potential Anal.*, 5(4): 403–415, 1996.

[9] P. Hajłasz: Sobolev spaces on metric-measure spaces. In *Heat kernels and analysis on manifolds, graphs, and metric spaces (Paris, 2002)*, vol. 338 of *Contemp. Math.*, 173–218. Amer. Math. Soc., Providence, RI, 2003.

[10] P. Hajłasz and P. Koskela: *Sobolev met Poincaré. Mem. Amer. Math. Soc.*, 145(688):x+101, 2000.

[11] J. Heinonen: *Lectures on analysis on metric spaces*. Universitext. Springer-Verlag, New York, 2001.

[12] J. Heinonen, P. Koskela, N. Shanmugalingam, J. Tyson: *Sobolev spaces on metric measure spaces. An approach based on upper gradients*. New Mathematical Monographs, 27. Cambridge University Press, Cambridge, 2015.

[13] J. Hu: A note on Hajłasz-Sobolev spaces on fractals. *J. Math. Anal. Appl.*, 280(1): 91–101, 2003.

[14] J. Hu, Y. Ji and Z. Wen: Hajłasz-Sobolev type spaces and p-energy on the Sierpinski gasket. *Ann. Acad. Sci. Fenn. Math*, 30 (2005), no. 1, 99-111.

[15] J. Hutchinson: Fractals and Self Similarity. *Indiana Univ. Math. J.*, 30 (1981), no. 5, 713–747.

[16] X. Jiang, D. Yang and W. Yuan: The grand Lusin-area characterization of Hajłasz-Sobolev spaces and Triebel-Lizorkin spaces. *Math. Nachr.* 286 (2013), no. 7, 691-711.

[17] A. Jonsson, and H. Wallin: *Function spaces on subsets of* $\mathbb{R}^n$. Math. Rep. 2 (1984), no. 1, 1-221.

[18] J. Kigami: *Analysis on fractals.* Cambridge Tracts in Mathematics, 143. Cambridge University Press, Cambridge, 2001.

[19] P. Koskela and E. Saksman: Pointwise characterizations of Hardy-Sobolev functions. *Math. Res. Lett.* 15 (2008), no. 4, 727-744.

[20] P. Koskela, D. Yang and Y. Zhou: A characterization of Hajłasz-Sobolev and Triebel-Lizorkin spaces via grand Littlewood-Paley functions. *J. Funct. Anal.* 258 (2010), no. 8, 2637-2661.

[21] P. Koskela, D. Yang and Y. Zhou: Pointwise characterizations of Besov and Triebel-Lizorkin spaces and quasiconformal mappings. *Adv. Math.* 226 (2011), no. 4, 3579-3621.

[22] K. Pietruska-Pałuba and A. Stos: Poincaré inequality and Hajłasz-Sobolev spaces on nested fractals. *Studia Math.* 218 (2013), no. 1, 126.

[23] J. Rissanen: Wavelets on self-similar sets and the structure of the spaces $M^{1,p}(E,\mu)$. *Ann. Acad. Sci. Fenn. Math. Diss.*, 125 (2002), 46. Dissertation, University of Jyäskyä, Jyväskylä, 2002.

[24] N. Shanmugalingam: Newtonian spaces: an extension of Sobolev spaces to metric measure spaces. *Rev. Mat. Iberoamericana*, 16(2): 243–279, 2000.

[25] R. Strichartz: *Differential equations on fractals. A tutorial.* Princeton University Press, Princeton, NJ, 2006. .

[26] D. Yang: New characterizations of Hajłasz-Sobolev spaces on metric spaces. *Sci. China Ser.* A 46 (2003), no. 5, 675-689.

[27] X. Zhou: Absolutely continuous functions on compact and connected One-dimensional spaces. *Preprint*, 2015.

[28] Y. Zhou: Criteria for optimal global integrability of Hajłasz-Sobolev functions, *Illinois J. Math.* 55 (2011), no. 3, 1083-1103 (2013).

Xiaodan Zhou, Department of Mathematics, University of Pittsburgh, Pittsburgh, PA 15260, USA, xiz78@pitt.edu